

HAEMOLYTIC URAEMIC SYNDROME ASSOCIATED WITH A FAMILY CLUSTER OF ENTEROHAEMORRHAGIC *ESCHERICHIA COLI*

Jeffrey N Hanna, Jan L Humphreys, Sian E Ashton, Denise M Murphy

Introduction

In early 1995 an outbreak of 23 cases of haemolytic uraemic syndrome (HUS) occurred in children (ranging from four months to 12 years of age) in South Australia.¹ Twenty of the cases were managed in a tertiary paediatric hospital in Adelaide, where 18 (90%) required dialysis.² A 4-year-old died and 12 months after discharge 5 of the surviving children still had significantly impaired renal function.²

In Australia HUS is usually caused by a subgroup of Shiga toxin-producing *Escherichia coli* known as enterohaemorrhagic *E. coli* (EHEC). The Shiga toxins cause cell damage and trigger an inflammatory process which initiates intravascular coagulation resulting in microthrombi forming in small blood vessels in the gut and kidney.³ The natural reservoir of EHEC is the gut of animals, particularly cattle and sheep. Hence HUS can be caused by contact with animal faeces, either directly or via contaminated, inadequately cooked food, particularly meat and dairy products. Most of the cases in the 1995 outbreak of HUS in South Australia had consumed (in the week before the onset of illness) an uncooked fermented sausage manufactured in Adelaide.¹ Subsequent molecular studies revealed an identical EHEC in both faeces of the cases and samples of the sausage.⁴

Because EHEC infection, and therefore HUS, can be foodborne, it is of considerable public health concern. Following the South Australian outbreak, HUS became a notifiable disease in Queensland in mid-1996, and EHEC in mid-2001. However, a recent case of HUS in north Queensland has identified several shortcomings in the management and investigation of HUS and EHEC infections; some of these shortcomings were also identified in a previous cluster of HUS cases that occurred in north Queensland in 2004.⁵

The HUS case and subsequent investigations

In early January 2007, the Tropical Population Health Network (TPHN) was notified by an infection control practitioner that a 14-month-old Caucasian girl had been hospitalised the previous day with HUS; she had become unwell four days

before being hospitalised. *Salmonella* Virchow was isolated from a diarrhoeal stool sample collected two days prior to her being hospitalised and from a stool sample collected on the day of admission.

The attending physician initially believed that the *Salmonella* infection was the cause of the HUS, and this led to problems in getting the (diarrhoeal) samples to the Queensland Health Scientific Services reference laboratory for screening for Shiga toxin (stx1 and stx2) gene (and therefore for EHEC). The initial sample was not forwarded, and the second sample was not forwarded frozen to the laboratory. When the latter sample was eventually screened stx genes were not detected. Therefore EHEC was never detected in the HUS case.

The child's parents were interviewed using the relevant OzFoodNet questionnaire; this did not reveal any suspect food items in the child's diet. However, it did reveal that the child and her two siblings had visited several commercial animal sanctuaries during the exposure period. At two of these the children had had direct contact with marsupials (particularly kangaroos and koalas) and apparently also with faeces from these animals. The parents stated that the children had no contact with other mammals at these sanctuaries, and no apparent contact with any bovine animals during the exposure period.

The child's two siblings attended a local child-care centre. Even though they were apparently asymptomatic, stool samples were collected (in mid-January) from both and the parents were requested by TPHN to keep them out of child care until the results of the stool tests were known.^{6,7} The child's twin sibling's stool was positive for the stx2 (but not the stx1) gene and the eaeA gene (which encodes a virulence factor: intimin) upon screening, and was culture positive for *E. coli* O55:H80, S. Aberdeen and S. Chailey. The child's 3-year-old brother's stool was also positive for the stx2 (but not the stx1) gene and the eaeA gene upon screening, and was culture positive for *E. coli* O55:HR. ('R' indicates that the organism had become rough in sub-cultures; once an EHEC becomes rough, the H antigen cannot be typed.) Both parents then had stool samples collected, but neither had evidence of EHEC upon screening.

The two siblings were voluntarily excluded from child-care until they were clear of the Shiga toxin in weekly stool samples. The 3-year-old and the twin sibling were able to return to child-care (after having two successive stool samples collected at least 48 hours apart clear of any evidence of Shiga toxin or EHEC⁷) 3.5 and 4.5 weeks, respectively, after having been first identified as being infected with EHEC. This delay created considerable difficulties for the parents, and repeated explanations of the importance of their exclusion from child-care were necessary.

The two siblings (and presumably the case) were infected with Shiga toxin-producing *E. coli* (all presumably with O55:H80), and the twins were infected with three different *Salmonella* serovars. This array of pathogens supported the hypothesis, as suggested from the parent interview, that the EHEC was acquired via animal contact rather than via a particular food item. For this reason, an assessment of the facilities and signs at the two animal sanctuaries was undertaken (about 2.5 weeks after the onset of the HUS) by environmental health officers. The public was encouraged to handle animals at both sanctuaries but there were no signs recommending hand washing after handling the animals at either sanctuary. At one sanctuary there were no hand washing facilities near the animal-handling areas. There were several food outlets in close proximity to the animal facilities.

It appeared that the management of the sanctuaries had little understanding of the potential infectious hazards associated with such facilities, and were uncertain of their responsibilities to minimise the risk of such hazards. Indeed, there are no guidelines in Queensland on how to minimise these risks for managers of commercial facilities that encourage the public to handle animals.

There was no evidence of Shiga toxin in faecal samples from koalas and kangaroos at either of the facilities, however, the samples were collected about a month after the onset of the HUS.

Discussion

This report describes three siblings infected with a Shiga toxin-producing EHEC; two who remained asymptomatic (presumably both with *E. coli* O55:H80), and their sibling (presumably infected with the same EHEC) who developed HUS. Both the *sxt2* and *eaeA* genes were detected in this EHEC; this combination of genes appears to be an important predictor of HUS.⁸

This is the second cluster of EHEC with HUS in north Queensland in three years.⁵ The two clusters have identified several issues of concern (Box).

Salmonella infections do not cause HUS,⁹ and the isolation of salmonellae from faecal samples from a HUS patient must be regarded as coincidental to the HUS. Screening for Shiga toxin and other virulence genes was undertaken on the *S. Virchow* isolated from the HUS child to prove this point to the attending physician; none of the genes were detected. Faecal samples from HUS cases must be forwarded promptly to a reference laboratory for screening for EHEC regardless of the isolation of *Salmonella* from the samples. Failure to do this may result in EHEC not being isolated from a case (as happened with the child with HUS in this cluster), and could impede the necessary investigations.

Issues of public health significance revealed by this family cluster of EHEC infections

- HUS is a notifiable disease. As it is a syndrome, notification has to come from clinicians: not only paediatricians but also haematologists, nephrologists, infectious diseases and intensive care physicians need to be aware of their responsibility to notify cases of HUS.
- As soon as HUS is diagnosed, a stool sample should be sent to the relevant reference laboratory for screening tests for Shiga toxins and EHEC.
- *Salmonella* infections do not cause HUS. Isolation of *Salmonella* from the stool of an HUS case must be considered as coincidental to the HUS.
- Stool samples being submitted for investigation for Shiga toxin and EHEC must be frozen soon after collection, and transported frozen to the reference laboratory.
- Children less than 5 years of age who attend child-care, and who are household contacts of a case of symptomatic EHEC infection, should be screened for EHEC even if they are asymptomatic. They should be excluded from child-care until two stool samples, collected at least 48 hours apart, are shown to be clear of the EHEC.
- Infectious disease hazards are associated with contact with animals in public facilities – ‘petting zoos’ – and the management of such facilities need to take measures to reduce the risk of these hazards.
- Guidelines on how these infectious hazards can be minimised need to be formally endorsed by the relevant agencies so that the management of petting zoos and other similar facilities can be made aware of their responsibilities.

The infecting dose of HUS is very low, and person-to-person transmission of EHEC is well documented, with transmission occurring among young children within families and in child care facilities.¹⁰ For this reason it is essential to screen the young siblings of EHEC HUS cases for the organism even if they are asymptomatic, and older siblings (and other close contacts) if they have any relevant symptoms.⁵ These individuals should be excluded from child care (or any workplace of concern) while the screening takes place, and may need to be further excluded should the screening indicate an EHEC infection.^{6,7}

There does not appear to be any published information as to whether marsupials act as reservoirs of EHEC. However, it is well recognised that macropods (kangaroos, wallabies) can be infected with salmonellae,¹¹ and an outbreak of human salmonellosis associated with contact with wallabies in a petting zoo has been reported from the United States of America.¹² Several other zoonotic (mostly enteric) infections have occurred following handling animals in petting zoos.^{12,13}

It is important that guidelines on how to minimise the risk of transmission of infections through handling animals be made readily available to those facilities that encourage the public to handle animals on-site. These guidelines should include educating the public; appropriate signage; providing hand washing facilities; ensuring adequate supervision of children; discouraging eating in animal contact areas; ensuring sick animals are not handled by the public; providing appropriate cleaning and infection of the animal holding area and ensuring the safe disposal of animal faeces.¹³ Such guidelines are available in several countries,^{14,15} and in South Australia;¹⁶ these guidelines are being used as templates for the drafting, currently in progress, of petting zoo guidelines for use in Queensland.

Acknowledgements

We wish to thank the Microbiological Diagnostic Unit PHL, Department of Microbiology and Immunology, University of Melbourne for the H antigen typing and confirmation of the O typing.

Author details

Jeffrey N Hanna, Public Health Physician¹

Jan L Humphreys, Public Health Nursing Officer²

Sian E Ashton, Environmental Health Officer¹

Denise M Murphy, Supervising Scientist³

1. Tropical Population Health Network, Queensland Health, Cairns, Queensland
2. Tropical Population Health Network, Queensland Health, Townsville, Queensland
3. Public Health Microbiology, Queensland Health Scientific Services, Coopers Plains, Queensland

Corresponding author: Dr J Hanna, Tropical Population Health Network, PO Box 1103, CAIRNS QLD 4870. Telephone: +61 7 4050 3604. Facsimile: +61 7 4031 1440. Email: Jeffrey_hanna@health.qld.gov.au

References

1. Centers for Disease Control and Prevention. Community outbreak of hemolytic uremic syndrome attributable to *Escherichia coli* O111:NM — South Australia, 1995. *MMWR Morb Mortal Wkly Rep* 1995;44:550–551, 557–558.
2. Henning PH, Tham EB, Martin AA, Beare TH, Jureidini KF. Haemolytic-uraemic syndrome outbreak caused by *Escherichia coli* O111:H-: clinical outcomes. *Med J Aust* 1998;168:552–555.
3. Elliott EJ, Robins-Browne RM. Hemolytic uremic syndrome. *Curr Probl Pediatr Adolesc Health Care* 2005;35:310–330.
4. Paton AW, Ratcliffe RM, Doyle RM, Seymour-Murray J, Davos D, Lanser JA, et al. Molecular microbiological investigation of an outbreak of hemolytic-uremic syndrome caused by dry fermented sausage contaminated with Shiga-like toxin-producing *Escherichia coli*. *J Clin Microbiol* 1996;34:1622–1627.
5. Morgan AK, Piispanen J, Humphreys J, Murphy D. A cluster of cases of haemolytic uraemic syndrome in north Queensland associated with a novel Shiga-like toxin-producing *Escherichia coli*. *Commun Dis Intell* 2005;29:193–196.
6. Guidelines for the control of infection with vero cytotoxin producing *Escherichia coli* (VTEC). Subcommittee of the PHL Advisory Committee on Gastrointestinal Infections. *Commun Dis Public Health* 2000;3:14–23.
7. Queensland Health. Enterohaemorrhagic *Escherichia coli* (EHEC) infection. In: *Queensland Health Guidelines for the Control of Communicable Diseases in the Community*. 3rd edn. Brisbane, Queensland Government, 2005.
8. Ethelberg S, Olsen KE, Scheutz F, Jensen C, Schiellerup P, Engberg J, et al. Virulence factors for hemolytic uremic syndrome, Denmark. *Emerg Infect Dis* 2004;10:842–847.
9. Flores FX, Jabs K, Thorne GM, Jaeger J, Linshaw MA, Somers MJ. Immune response to *Escherichia coli* O157: H7 in hemolytic uremic syndrome following salmonellosis. *Pediatr Nephrol* 1997;11:488–490.
10. Ludwig K, Sarkim V, Bitzan M, Karmali MA, Bobrowski C, Ruder H, et al. Shiga toxin-producing *Escherichia coli* infection and antibodies against stx1 and stx2 in household contacts of children with enteropathic hemolytic-uremic syndrome. *J Clin Microbiol* 2002;40:1773–1782.
11. Thomas AD, Forbes-Faulkner JC, Speare R, Murray C. Salmonellosis in wildlife from Queensland. *J Wildl Dis* 2001;37:229–238.
12. Steinmuller N, Demma L, Bender JB, Eidson M, Angulo FJ. Outbreaks of enteric disease associated with animal contact: not just a foodborne problem anymore. *Clin Infect Dis* 2006;43:1596–1602.
13. Bender JB, Shulman SA, Animals in Public Contact subcommittee; National Association of State Public Health Veterinarians. Reports of zoonotic disease outbreaks associated with animal exhibits and availability of recommendations for preventing zoonotic disease transmission from animals to people in such settings. *J Am Vet Med Assoc* 2004;224:1105–1109.

14. Health and Safety Executive. Avoiding ill health at open farms – Advice to farmers (with teachers' supplement). Agriculture Information Sheet no. 23 (revised). June 2000. Available from: <http://www.hse.gov.uk/pubns/ais23.pdf> Accessed February 2007.
15. National Association of State Public Health Veterinarians, Inc. (NASPHV). Compendium of measures to prevent disease associated with animals in public settings, 2005. *MMWR Recomm Rep* 2005;54 RR-4:1–13.
16. Communicable Disease Control Branch and Environmental Health Branch. Petting zoo infection control guideline for petting zoo operators, education and childcare services and environmental health officers. Department of Human Services. February 2002. Available from: <http://www.dh.sa.gov.au/pehs/PDF-files/petting-zoos-guidelines.pdf> Accessed on March 2007.

CURRENT DEVELOPMENTS IN VARICELLA-ZOSTER VIRUS DISEASE PREVENTION

A report on the varicella-zoster virus workshop convened by the National Centre for Immunisation Research and Surveillance of Vaccine Preventable Diseases on 16–17 November 2006

Anita E Heywood, Kristine K Macartney, C Raina MacIntyre, Peter B McIntyre

Introduction

On 16 and 17 November 2006, the National Centre for Immunisation Research and Surveillance of Vaccine Preventable Diseases (NCIRS) hosted a workshop on varicella-zoster virus (VZV) disease. The workshop was aimed at presenting the latest information on the clinical, epidemiological, and diagnostic aspects of both primary varicella ('chickenpox') and herpes zoster (HZ or 'shingles') both in Australia and internationally, and to highlight important developments in the prevention of these diseases by vaccination. This workshop was held at a significant stage in the control of VZV disease in Australia with the recent addition of the varicella vaccine to the National Immunisation Program (NIP) schedule, the anticipated availability of combination measles-mumps-rubella-varicella (MMRV) vaccines for use in children, and the availability of a zoster vaccine for use in older adults to prevent reactivation of VZV causing HZ.

The workshop was attended by prominent international researchers and leading Australian experts. All state and territory jurisdictions were represented and participated in panel discussions, particularly with the regard to disease surveillance. The first day of the workshop was devoted to varicella disease with presentations on the clinical features, current epidemiology, the Australian varicella vaccination program, and the impact of varicella vaccination in the United States of America (USA). An overview of the development of the MMRV vaccines was provided, and the day closed with a panel discussion of the issues surrounding varicella vaccine scheduling.

The second day focused on HZ with presentations on the burden of disease in Australia, the pathologic mechanisms and diagnostics. An overview and update on data from the zoster vaccine clinical trials was presented. The day concluded with state and territory representatives presenting plans for disease surveillance, and a panel discussion focusing on the best approach for the control of VZV disease in Australia.

Day one – varicella

Clinical overview

Professor Margaret Burgess, NCIRS, began proceedings with a presentation on clinical features of primary VZV disease including varicella (chickenpox), and neonatal and congenital varicella. As she highlighted, varicella is usually a relatively mild disease of childhood, however, complications (such as pneumonia, secondary bacterial infections and neurologic conditions) occur in approximately 1% of cases, especially those most at risk such as neonates, the immunosuppressed, pregnant women, adolescents, adults and those with pre-existing co-morbidities.^{1,2} Professor Burgess presented the results of community-based surveys and seroprevalence studies in Australia that indicate that the majority of the burden of varicella is in childhood and adolescence with almost 90% of cases occurring before the age of 20 years and the most common age of acquisition between 5–9 years of age.³

Congenital and neonatal varicella are rare in Australia with the Australian Paediatric Surveillance Unit (APSU) reporting 44 cases of neonatal varicella